

**PhD course in Basic Biostatistics - Day 7**  
 Erik Parner, Department of Biostatistics, Aarhus University©

Post term delivery and parity  
 A two by two table

Odds and odds ratios

- Odds ratio and the relative risk
- Estimating the odds ratio
- Relative measures - why inference on the log-scale?
- Odds ratios, Confidence Intervals and testing

Odds ratio and logistic regression

Post term delivery and age of the woman

- A simple logistic regression model
- Understanding the model and the parameters
- The estimates
- Formulations
- The risk versus age curves

Erik Parner Basic Biostatistics - Day 7

1

Post term delivery and parity - adjusting for age

Two simple logistic regression models  
 Comparing the slopes

A logistic regression model with the same slope

Understanding the parameters

The estimates

Formulation

A risk versus age and parity plot

Why use logistic regression?

Linear and logistic regression models - a comparison

Why do we need regression models

- Adjustment
- Effect modification
- Prediction

Erik Parner Basic Biostatistics - Day 7

2

**Example: Post term delivery and parity**

**Question:** How does the risk of post term delivery depend on parity?

**Data:** Parity and gestational age for 12,311 women in the age of 20 to 39. Post term delivery defined as a gestational age larger than 40 weeks.

Parity	N	Postterm	Risk
First child	5,938	1,722	29.0 (28.8; 30.2)%
Not first child	6,373	1,677	26.3 (25.2; 27.4)%
Total	12,311	3,399	27.6 (26.8; 28.4)%

**Model:** Independent samples from two binomial distributions.

Let  $\pi_0$  and  $\pi_1$  be the probability (risk) of post term delivery among women giving birth to their first child or not, respectively.

Erik Parner

Basic Biostatistics - Day 7

3

**Example: Post term delivery and parity**

The assumptions behind the model was discussed on day 4.

On that day we also looked at three different measures of associations: Risk Difference, Relative Risk and Odds Ratio. And the chi-squared test for no association.

Today we will look closer at the Odds Ratio.

Risk difference	-2.7 (-4.3; -1.1)%
Relative risk	0.91 (0.86; 0.96)
Odds ratio	0.87 (0.81; 0.95)
$\chi^2=11.09$	$p=0.001$

In the table above we compare  $\pi_1$  to  $\pi_0$ , i.e. women giving birth to their first child is the reference group.

We see that the risk is (statistically significant) smaller if the woman already had a child.

Erik Parner

Basic Biostatistics - Day 7

4

### Odds and risk

The **odds** is defined as  $\pi/(1-\pi)$ , i.e. the probability of post term delivery divided by the probability of not having a post term delivery.

$$odds = \frac{\pi}{1-\pi}$$

If the odds is equal to 0.5=1/2, then the risk of post term delivery is only **half** of the risk of not having a post term delivery.

We can also go from odds to risk:

$$\pi = \frac{odds}{1+odds}$$

We see that

$$odds = 0.5 \quad \text{gives} \quad \pi = 0.5/(1+0.5)=0.3333.$$

$$odds = 1 \quad \text{gives} \quad \pi = 1/(1+1)=0.5.$$

Erik Parner

Basic Biostatistics - Day 7

5

### Odds and odds ratios

The odds ratio comparing parity>0 to the reference is given by

$$OR_{10} = \frac{odds_1}{odds_0} = \frac{\pi_1/(1-\pi_1)}{\pi_0/(1-\pi_0)} = \frac{\pi_1 \cdot (1-\pi_0)}{\pi_0 \cdot (1-\pi_1)}$$

It is easily seen that  $\pi_1 = \pi_0 \Leftrightarrow odds_1 = odds_0 \Leftrightarrow OR = 1$

OR has **nice properties**:

Switching reference group or event will just lead to 1/OR, e.g.

$$OR_{01} = \frac{odds_0}{odds_1} = \frac{1}{OR_{10}}$$

And of course the estimates and confidence intervals will transform similarly.

$$OR_{01} : \frac{1}{0.87} \left( \frac{1}{0.95}; \frac{1}{0.81} \right) = 1.14(1.06;1.24)$$

Erik Parner

Basic Biostatistics - Day 7

6

### Odds ratios and relative risks

The odds ratio is related to the relative risk:

$$OR_{10} = \frac{\pi_1 \cdot (1-\pi_0)}{\pi_0 \cdot (1-\pi_1)} = RR_{10} \cdot \frac{(1-\pi_0)}{(1-\pi_1)}$$

We can see that if **the event is rare**, i.e. both  $\pi_1$  and  $\pi_0$  are small, then the last ratio is close to 1/1=1.

So for a **rare event** we have:

$$OR \approx RR$$

Erik Parner

Basic Biostatistics - Day 7

7

### Estimating the odds ratios

The odds ratio is of course estimated by:

$$\widehat{OR}_{10} = \frac{\widehat{\pi}_1 \cdot (1-\widehat{\pi}_0)}{\widehat{\pi}_0 \cdot (1-\widehat{\pi}_1)}$$

Another way to find the estimate is to make the 'classical' 2x2 table:

Exposed	Event	
	Yes	No
Yes	a	b
No	c	d

$$\widehat{OR}_{10} = \frac{a \cdot d}{b \cdot c}$$

Parity>0	Post term	
	Yes	No
Yes	1,677	4,696
No	1,722	4,216

$$\widehat{OR}_{10} = \frac{1,677 \cdot 4,216}{1,722 \cdot 4,696} = 0.8743$$

Erik Parner

Basic Biostatistics - Day 7


8

### Odds ratios - why inference on the log-scale

The odds ratio is limited to be positive.

A value in the interval **0 to 1** corresponds to lower risk among the Parity>0.

A value from **1 to infinity** corresponds to higher risk among the Parity>0



If we switch "exposed" and "unexposed" we get



Erik Parner

Basic Biostatistics - Day 7


9

### Odds ratios - why inference on the log-scale


The **log-odds ratio** is not limited.

A value in the interval **-infinity to 0** corresponds to lower risk among the Parity>0.

A value from **0 to infinity** corresponds to higher risk among the Parity>0



If we switch "exposed" and "unexposed" we get



**Symmetry on the log scale!!!**

Erik Parner

Basic Biostatistics - Day 7

10

### Odds ratios - Approx. CI (Woolf/Wald)

So on the **log scale** we have a symmetric measure of association.

On the **log scale** it makes sense to find the CI as 'usual', i.e. as **estimate  $\pm 1.96 \cdot se$** .

Using the notation from page 8 we have:

$$se\left(\ln\left(\widehat{OR}\right)\right) = \sqrt{\frac{1}{a} + \frac{1}{b} + \frac{1}{c} + \frac{1}{d}}$$

$$se\left(\ln\left(\widehat{OR}\right)\right) = \sqrt{\frac{1}{1,677} + \frac{1}{4,696} + \frac{1}{1,711} + \frac{1}{4,216}} = 0.0403$$

$$CI \ln(OR) = \ln(0.8743) \pm 1.96 \cdot 0.0403 = (-0.2134; -0.0552)$$

$$CI OR = (\exp(-0.2134); \exp(-0.0552)) = (0.81; 0.95)$$

Erik Parner

Basic Biostatistics - Day 7

11

### Odds ratios - Testing

If one wants to test a hypothesis that the odds ratio has a specific value:  $OR = OR_0$ ,

then this is also done on the log-scale:

$$z_{obs} = \frac{\ln\left(\widehat{OR}\right) - \ln\left(OR_0\right)}{se\left(\ln\left(\widehat{OR}\right)\right)}$$

Could the odds be reduced by 10%, i.e.  $H: OR=0.9$ ?

$$z_{obs} = \frac{\ln(0.8743) - \ln(0.9)}{0.0403} = -0.719$$

$$p = 2 \cdot \Pr(z > |z_{obs}|) = 2 \cdot \Pr(z < -|z_{obs}|) = 0.47$$

The hypothesis cannot be rejected.

```
disp ( ln(0.8743) - ln(0.9) ) / 0.0403
disp 2*normal( abs(-0.719) )
```

Erik Parner

Basic Biostatistics - Day 7

12

### Odds ratios - logistic regression

Here we will see how one can find the odds ratio by logistic regression.

Let  $Par1$  be an indicator variable for Parity>0, i.e.  $Par1=1$  if parity>0 and  $Par1=0$  if Parity==0.

Now we will look at the (logistic regression) model:

$$\log(odds) = \beta_0 + \beta_1 \cdot Par1$$

This is equivalent to:

$$odds = \exp(\beta_0 + \beta_1 \cdot Par1) = \exp(\beta_0) \cdot \exp(\beta_1)^{Par1}$$

and

$$\pi = \frac{\exp(\beta_0 + \beta_1 \cdot Par1)}{1 + \exp(\beta_0 + \beta_1 \cdot Par1)}$$

Erik Parner

Basic Biostatistics - Day 7

13

### Odds ratios - logistic regression

$$\log(odds) = \beta_0 + \beta_1 \cdot Par1 \quad odds = \exp(\beta_0) \cdot \exp(\beta_1)^{Par1}$$

We see that if  $Parity=0$  then we have:

$$\log(odds) = \beta_0 \quad odds = \exp(\beta_0)$$

and if  $Parity>0$  then we have

$$\log(odds) = \beta_0 + \beta_1 \quad odds = \exp(\beta_0) \cdot \exp(\beta_1)$$

Combining we have

$$OR_{10} = \frac{odds(\text{if parity}>0)}{odds(\text{if parity}=0)} = \frac{\exp(\beta_0) \cdot \exp(\beta_1)}{\exp(\beta_0)} = \exp(\beta_1)$$

Erik Parner

Basic Biostatistics - Day 7

14

### Odds ratios - logistic regression

$$\log(odds) = \beta_0 + \beta_1 \cdot Par1 \quad odds = \exp(\beta_0) \cdot \exp(\beta_1)^{Par1}$$

In summary we have that in the model:

The "intercept"  $\beta_0$  is the log odds in the "reference group".

The "slope"  $\beta_1$  is the log OR.

That is, we can find the odds ratio from before by what is called a logistic regression model.

So the computer will give us estimates and confidence intervals for the odds in the reference group and the odds ratio comparing the 'exposed' to the reference.

Erik Parner

Basic Biostatistics - Day 7

15

### Post term delivery and age

We know that the age distribution among the two groups of women is different - the women giving birth for the first time are on average younger (27.5 years versus 30.2 years in the data set).

It might be relevant to compare the two groups after "adjustment for age".

We will start by modeling the association between post term delivery and age among the women with  $Parity=0$ .

The simplest logistic regression model is:

$$\log(odds) = \alpha_0 + \alpha_1 \cdot Age \quad odds = \exp(\alpha_0) \cdot \exp(\alpha_1)^{Age}$$

To get a sensible reference age:

$$\log(odds) = \alpha_0 + \alpha_1 \cdot (Age - 30) \quad odds = \exp(\alpha_0) \cdot \exp(\alpha_1)^{Age-30}$$

Erik Parner

Basic Biostatistics - Day 7

16

Post term delivery and age (Parity==0)								
$\log(\text{odds}) = \alpha_0 + \alpha_1 \cdot (\text{Age} - 30)$			$\text{odds} = \exp(\alpha_0) \cdot \exp(\alpha_1)^{\text{Age}-30}$					
$\text{Age} = 30: \log(\text{odds}) = \alpha_0$			$\text{odds} = \exp(\alpha_0)$					
$\text{Age} = 31: \log(\text{odds}) = \alpha_0 + \alpha_1$			$\text{odds} = \exp(\alpha_0) \cdot \exp(\alpha_1)$					
$\text{Age} = 20: \log(\text{odds}) = \alpha_0 - 10 \cdot \alpha_1$			$\text{odds} = \exp(\alpha_0) \cdot \exp(\alpha_1)^{-10}$					
$\text{Age} = 21: \log(\text{odds}) = \alpha_0 - 9 \cdot \alpha_1$			$\text{odds} = \exp(\alpha_0) \cdot \exp(\alpha_1)^{-9}$					
$\text{Age} = 25: \log(\text{odds}) = \alpha_0 - 5 \cdot \alpha_1$			$\text{odds} = \exp(\alpha_0) \cdot \exp(\alpha_1)^{-5}$					
$OR_{31 \text{ vs } 30} = \exp(\alpha_0) \cdot \exp(\alpha_1) / \exp(\alpha_0) = \exp(\alpha_1)$								
$OR_{21 \text{ vs } 20} = \exp(\alpha_0) \cdot \exp(\alpha_1)^{-9} / (\exp(\alpha_0) \cdot \exp(\alpha_1)^{-10}) = \exp(\alpha_1)$								
$OR_{25 \text{ vs } 20} = \exp(\alpha_0) \cdot \exp(\alpha_1)^{-5} / (\exp(\alpha_0) \cdot \exp(\alpha_1)^{-10}) = \exp(\alpha_1)^5$								

Erik Parner

Basic Biostatistics - Day 7

17

Post term delivery and age (Parity==0)								
$\log(\text{odds}) = \alpha_0 + \alpha_1 \cdot (\text{Age} - 30)$			$\text{odds} = \exp(\alpha_0) \cdot \exp(\alpha_1)^{\text{Age}-30}$					
We saw: $\text{Age} = 30: \text{odds} = \exp(\alpha_0)$								
$OR_{31 \text{ vs } 30} = \exp(\alpha_1)$								
$OR_{21 \text{ vs } 20} = \exp(\alpha_1)$								
$OR_{25 \text{ vs } 20} = \exp(\alpha_1)^5$								
That is, $\exp(\alpha_0)$ is the odds in the reference ( $\text{Age}=30$ ) $\exp(\alpha_1)$ is the <b>OR</b> for 1 year difference. and								
$OR_{5 \text{ years}} = OR_{1 \text{ year}}^5$								

Erik Parner

Basic Biostatistics - Day 7

18

Post term delivery and age (Parity==0)																													
Using a computer we get:																													
<table border="1"> <thead> <tr> <th colspan="6">log odds scale</th> </tr> <tr> <th></th><th>est</th><th>se</th><th>lower</th><th>upper</th><th>H = 0</th></tr> </thead> <tbody> <tr> <td>Const</td><td>-0.8446</td><td>0.0332</td><td>-0.9096</td><td>-0.7795</td><td>-25.44 &lt;0.0001</td></tr> <tr> <td>Age-30</td><td>0.0207</td><td>0.0071</td><td>0.0069</td><td>0.0345</td><td>2.93 0.003</td></tr> </tbody> </table>						log odds scale							est	se	lower	upper	H = 0	Const	-0.8446	0.0332	-0.9096	-0.7795	-25.44 <0.0001	Age-30	0.0207	0.0071	0.0069	0.0345	2.93 0.003
log odds scale																													
	est	se	lower	upper	H = 0																								
Const	-0.8446	0.0332	-0.9096	-0.7795	-25.44 <0.0001																								
Age-30	0.0207	0.0071	0.0069	0.0345	2.93 0.003																								
<table border="1"> <thead> <tr> <th colspan="6">Exp odds scale</th> </tr> <tr> <th></th><th>est</th><th>se</th><th>lower</th><th>upper</th><th>H = 1</th></tr> </thead> <tbody> <tr> <td>Const</td><td>0.4297</td><td>0.4027</td><td>0.4586</td><td>-25.44</td><td>&lt;0.0001</td></tr> <tr> <td>Age-30</td><td>1.0209</td><td>1.0069</td><td>1.0351</td><td>2.93</td><td>0.003</td></tr> </tbody> </table>						Exp odds scale							est	se	lower	upper	H = 1	Const	0.4297	0.4027	0.4586	-25.44	<0.0001	Age-30	1.0209	1.0069	1.0351	2.93	0.003
Exp odds scale																													
	est	se	lower	upper	H = 1																								
Const	0.4297	0.4027	0.4586	-25.44	<0.0001																								
Age-30	1.0209	1.0069	1.0351	2.93	0.003																								
Note, only the estimates and the confidence intervals should be transformed!																													

Erik Parner

Basic Biostatistics - Day 7

19

Post term delivery and age (Parity==0)																													
<table border="1"> <thead> <tr> <th colspan="6">CI</th></tr> <tr> <th></th><th>est</th><th>lower</th><th>upper</th><th>z</th><th>p</th></tr> </thead> <tbody> <tr> <td>Odds if Age=30</td><td>0.4297</td><td>0.4027</td><td>0.4586</td><td>-25.44</td><td>&lt;0.0001</td></tr> <tr> <td>One years age dif.</td><td>1.0209</td><td>1.0069</td><td>1.0351</td><td>2.93</td><td>0.003</td></tr> </tbody> </table>						CI							est	lower	upper	z	p	Odds if Age=30	0.4297	0.4027	0.4586	-25.44	<0.0001	One years age dif.	1.0209	1.0069	1.0351	2.93	0.003
CI																													
	est	lower	upper	z	p																								
Odds if Age=30	0.4297	0.4027	0.4586	-25.44	<0.0001																								
One years age dif.	1.0209	1.0069	1.0351	2.93	0.003																								
From odds to probability:																													
$\Pr(\text{post term if Age}=30) = \frac{0.4297}{1+0.4297} \left( \frac{0.4027}{1+0.4027}; \frac{0.4586}{1+0.4586} \right)$ $= 30.1(28.7;31.4)\%$																													
Five years age difference:																													
$OR_{5 \text{ years}} = 1.0209^5 (1.0069; 1.0351)^5$ $= 1.11(1.03; 1.19)$																													

Erik Parner

Basic Biostatistics - Day 7

20

### Post term delivery and age (Parity==0) - Formulations

#### Methods

The risk of post term delivery among women giving birth for the first time was described by a logistic regression model with age as a continuous variable. ...

#### Results

We found that five year age difference corresponds to an odds ratio of 1.11(1.03; 1.19). A 30 year old woman giving birth for the first time has 30(29;31)% risk of post term delivery.

#### Conclusion

The risk of post delivery among women giving birth for the first time increases with the age of the woman....

Erik Parner

Basic Biostatistics - Day 7

21

### Logistic regression checking the model

It is outside the scope of this course to go into details on how to check the model, so we will just state the assumptions behind the model:

1. All the observations should be **independent**.
2. There is exactly the same **two possible outcomes** for each observation.
3. The log odds is a **linear function** of age.

The last assumption can to some extend be checked by plotting the fitted regression line and the observed odds (with 95% CI) for each distinct age.


Erik Parner

Basic Biostatistics - Day 7

22

### Post term delivery and age (Parity==0)

$$\log(\text{odds}) = \alpha_0 + \alpha_1 \cdot (\text{Age} - 30)$$




Erik Parner

Basic Biostatistics - Day 7

23

### Post term delivery and age (Parity==0)

$$\text{risk} = \frac{\exp(\alpha_0 + \alpha_1 \cdot (\text{Age} - 30))}{1 + \exp(\alpha_0 + \alpha_1 \cdot (\text{Age} - 30))}$$



Not linear! But almost

Erik Parner


Basic Biostatistics - Day 7

24

### Post term delivery and age (Parity==0)

Extrapolating we see the non-linearity.

$$\text{risk} = \frac{\exp(\alpha_0 + \alpha_1 \cdot (\text{Age} - 30))}{1 + \exp(\alpha_0 + \alpha_1 \cdot (\text{Age} - 30))}$$



Erik Parner

Basic Biostatistics - Day 7

25

### Post term delivery parity adjusting for age

We now know that

- The risk of post term delivery **increases with age** (among women with **parity==0**).
- The risk of post term delivery is **smaller for Parity>0**.
- Women with **Parity>0** are older.

From this we can deduce that **adjusting for age** (if reasonable) will **increase** the difference between the two parity groups.

We now show how to find an age adjusted estimate, when we assume a linear "effect" of age on log odds.

Erik Parner

Basic Biostatistics - Day 7

26

### Post term delivery and age

We fit the same model to the **Parity>0** group and then look at the difference:

log odds	Slope				log odds Age==30			
	est	se	lower	upper	est	se	lower	upper
Parity>0	0.025	0.007	0.012	0.039	-1.037	0.029	-1.093	-0.981
Parity==0	0.021	0.007	0.007	0.035	-0.845	0.033	-0.910	-0.780
Difference	0.005	0.010	-0.015	0.024	-0.193	0.044	-0.279	-0.107

The **standard errors** of the differences are found as usual:

$$\text{se}(\text{est}_{\text{parity}>0} - \text{est}_{\text{parity}==0}) = \sqrt{\text{se}_{\text{parity}>0}^2 + \text{se}_{\text{parity}==0}^2}$$

We see that we can assume the slopes to be identical, we could also test the hypothesis:

$$z = \frac{0.005 - 0}{0.010} = 0.5 \quad p = 64\%$$

Erik Parner

Basic Biostatistics - Day 7

27

### Post term delivery and age - assuming identical slopes

If we **assume identical slopes**, then we can write the model:

$$\log(\text{odds}) = \gamma_0 + \gamma_1 \cdot (\text{Age} - 30) + \gamma_2 \cdot \text{Par1}$$

$$\text{odds} = \exp(\gamma_0) \cdot \exp(\gamma_1)^{(\text{Age}-30)} \cdot \exp(\gamma_2)^{\text{Par1}}$$

We see that

$\exp(\gamma_0)$  is the odds among 30-year old with **Parity==0**.

$\exp(\gamma_0) \cdot \exp(\gamma_2)$  is the odds among 30-year old with **Parity>0**.

$\exp(\gamma_1)^A$  is the odds ratio for the age difference **A** years among women in the same **Parity** group.

$\exp(\gamma_2)$  is the odds ratio comparing **Parity>0** to **Parity==0**, at the same age.

Erik Parner

Basic Biostatistics - Day 7

28

### Post term delivery and age - assuming identical slopes

The model is easily fitted by a computer:

log odds	Slope				log odds Age==30			
	est	se	lower	upper	est	se	lower	upper
Parity>0	0.023	0.005	0.013	0.033	-1.036	0.029	-1.092	-0.981
Parity==0					-0.839	0.031	-0.900	-0.778
Difference	0				-0.197	0.043	-0.281	-0.114

The age adjusted OR comparing Parity>0 to Parity==0:

$$OR_{10} : \exp(-0.197) \left( \exp(-0.281); \exp(-0.114) \right) = 0.821 (0.755; 0.892)$$

and if we compare Parity==0 to Parity>0:

$$OR_{01} : \frac{1}{0.821} \left( \frac{1}{0.892}; \frac{1}{0.755} \right) = 1.22 (1.12; 1.32)$$

Erik Parner

Basic Biostatistics - Day 7

29

### Post term delivery and Parity - Formulations

#### Methods

The risk of post term delivery among women was modeled by a logistic regression model with age as a continuous variable....

#### Results version1

Comparing Parity>0 to Parity==0 the crude odds ratio was 0.87(0.81;0.95). The age adjusted odds ratio was 0.82(0.76;0.89).

#### Results version2

Comparing Parity==0 to Parity>0 the crude odds ratio was 1.14(1.06;1.24). The age adjusted odds ratio was 1.22(1.12;1.32).

#### Conclusion ??

Women giving birth for the first time have a 22% (12%;32%) higher odds of post term delivery compared to other women of the same age. ....

Erik Parner

Basic Biostatistics - Day 7

30

The odds can be predicted by

$$odds = \exp(\gamma_0 + \gamma_1 \cdot (Age - 30) + \gamma_2 \cdot Par1)$$

The risk can be predicted by

$$risk = \frac{\exp(\gamma_0 + \gamma_1 \cdot (Age - 30) + \gamma_2 \cdot Par1)}{1 + \exp(\gamma_0 + \gamma_1 \cdot (Age - 30) + \gamma_2 \cdot Par1)}$$

Odds ratio is not always relative risk

A women of age 30 in the data set:

$$\begin{array}{ll} \text{Parity 0: odds=0.3547} & \text{risk=0.2618} \\ \text{Parity 1: odds=0.4321} & \text{risk=0.3017} \\ \text{OR=1.22} & \text{RR=1.15} \end{array}$$


Erik Parner

Basic Biostatistics - Day 7

31

### Post term delivery and age - assuming identical slopes

$$\log(odds) = \gamma_0 + \gamma_1 \cdot (Age - 30) + \gamma_2 \cdot Par1$$




Erik Parner

Basic Biostatistics - Day 7

32

**Post term delivery and age - assuming identical slopes**

$$risk = \frac{\exp(\gamma_0 + \gamma_1 \cdot (Age - 30) + \gamma_2 \cdot Par1)}{1 + \exp(\gamma_0 + \gamma_1 \cdot (Age - 30) + \gamma_2 \cdot Par1)}$$




Erik Parner

Basic Biostatistics - Day 7

33

**Post term delivery and age - assuming identical slopes**

$$risk = \frac{\exp(\gamma_0 + \gamma_1 \cdot (Age - 30) + \gamma_2 \cdot Par1)}{1 + \exp(\gamma_0 + \gamma_1 \cdot (Age - 30) + \gamma_2 \cdot Par1)}.$$



Erik Parner

Basic Biostatistics - Day 7

34

**Why use logistic regression**

There is a **long tradition** for using logistic regression when considering binary outcome. Some of the reasons are:

It is the **mathematical nicest** model for binary outcome, and hence the **first** type of models that was included in the statistical software.

If you have a **case-control** design, then you want to work with odds ratios.

If the **event is rare**, then it will give you **relative risk** estimates.

It is one of the few models for binary data that ensures that the estimated probability is **between zero and one**.

Erik Parner

Basic Biostatistics - Day 7

35

**Linear and logistic regression - a comparison**

In a **linear regression** the **outcome is continuous**: Lung function, Blood pressure, BMI, concentrations...

In a **logistic regression** the **outcome is binary**: Post term delivery, gender, dead/alive, sick/ well, BMI>30..

Neither of the models make **any assumptions** about the **explanatory variable**!!

In both models they can be continuous, binary or categorical.

In **both models** we have to assume **independence** between observations.

In **both models** we assume linearity - of expected value or the log odds.

Both models are readily fitted by standard statistical packages.

Erik Parner

Basic Biostatistics - Day 7

36

### Regression models in general - why? Adjustment

You have now looked at two of the most commonly used regression models in their most simple forms, involving one continuous and one binary explanatory variable.

You have seen how one can use such models for **adjustment**: What is the 'effect' of the binary 'exposure' when **adjusting** for the continuous variable?

Exactly the same models could answer the question: What is the 'effect' (slope) of the **continuous** 'exposure' when **adjusting** for the binary variable?  
E.g. what is the increase in risk of post term delivery associated with age when we **adjust** for parity?  
Often one has several explanatory variables, a mixture of continuous, binary and categorical and the purpose is to **adjust for more than one**.  
In such case one might apply a **multiple regression** model.

Erik Parner

Basic Biostatistics - Day 7

37

### Regression models in general - why? Effect modification

We have also seen how we could compare the 'effect' of one explanatory variable for subgroups described by another explanatory variable (**effect modification**):

What is the difference in the PEFR-height relationship for men and women?  
What is the difference in the Risk-age relationship for the two parity groups?

Typically by **comparing the slopes**.

Often one has several explanatory variables, a mixture of continuous, binary and categorical and the purpose is to **model effect modification** between explanatory variables.  
In such case one might apply a **multiple regression** model.

Erik Parner

Basic Biostatistics - Day 7

38

### Regression models in general - why? Prediction

We could also have used exactly the same models for **prediction/prognosis**:

What is the expected PEFR for a person with a given sex and a given height?  
What is the risk of post term delivery for women of a given age having her first child?  
Often one has several explanatory variables, a mixture of continuous, binary and categorical and the purpose is to make **prediction** for a person with a given set of characteristics.  
In such a case one might apply a **multiple regression** model.

Erik Parner

Basic Biostatistics - Day 7

39