


Multiple linear regression 2
Stata 11
Morten Frydenberg ©
Department of Biostatistics, Aarhus Univ, Denmark

Categorical variables in regression models.
The changing reference level
Interaction/effect modification
Interaction between a categorical and continuous variable
Interaction between two categorical variables

Morten Frydenberg Linear and Logistic regression - Note 2.2 1

Categorical variable in regression models

The age distribution:



Let us divide age into three agegroups ,
0: age≤40, 1: 40<age≤50, 2: 50<age
and consider the new model
$$\ln(sbp) = \alpha_0 + \alpha_1 \cdot age1 + \alpha_2 \cdot age2 + \alpha_3 \cdot woman + \alpha_4 \cdot \ln\left(\frac{bmi}{25}\right) + E$$

Morten Frydenberg Linear and Logistic regression - Note 2.2 2

Categorical variable : age group 0 reference

$$\ln(sbp) = \alpha_0 + \alpha_1 \cdot age1 + \alpha_2 \cdot age2 + \alpha_3 \cdot woman + \alpha_4 \cdot \ln\left(\frac{bmi}{25}\right) + E$$

age1 is one if a person is in age group 1 and zero otherwise
age2 is one if a person is in age group 2 and zero otherwise

The expected ln(sbp) in the three age groups will be:

age < 40: $\ln(sbp) = \alpha_0 + \alpha_3 \cdot woman + \alpha_4 \cdot \ln(bmi/25)$
40 ≤ age < 50: $\ln(sbp) = \alpha_0 + \alpha_1 + \alpha_3 \cdot woman + \alpha_4 \cdot \ln(bmi/25)$
50 ≤ age: $\ln(sbp) = \alpha_0 + \alpha_2 + \alpha_3 \cdot woman + \alpha_4 \cdot \ln(bmi/25)$

We see that α₁ is the adjusted difference in ln(sbp) when comparing a person in the second group with one in the first group.

And α₂ is the adjusted difference in ln(sbp) when comparing a person in the third group with one in the first group.

Morten Frydenberg Linear and Logistic regression - Note 2.2 3

Categorical variable : age group 0 reference

$$\ln(sbp) = \alpha_0 + \alpha_1 \cdot age1 + \alpha_2 \cdot age2 + \alpha_3 \cdot woman + \alpha_4 \cdot \ln\left(\frac{bmi}{25}\right) + E$$

Finally we see that α₀ is the expected ln(sbp) for a man in the first (reference) age group, with bmi=25.

In most programs the model is fitted by first generating the grouping variable and then making the regression telling the program which variables are categorical.

In Stata this done like this:

```
egen agegrp3=cut(age), at(0,40,50,120) label  
regress lnSBP woman i.agegrp3 lnBMI25
```

This is categorical

Morten Frydenberg Linear and Logistic regression - Note 2.2 4

Categorical variable : age group 0 reference

$$\ln(sbp) = \alpha_0 + \alpha_1 \cdot age1 + \alpha_2 \cdot age2 + \alpha_3 \cdot woman + \alpha_4 \cdot \ln\left(\frac{bmi}{25}\right) + E$$

OUTPUT:					
Source	SS	df	MS	Number of obs = 200	
Model	.980169926	4	.245042482	F(4, 195) = 11.20	
Residual	4.26524771	195	.021873065	Prob > F = 0.0000	
				R-squared = 0.1869	
				Adj R-squared = 0.1702	
Total	5.24541764	199	.026358883	Root MSE = .1479	

lnSBP	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
woman	.0035403	.0212026	0.17	0.868	-.0382757	.0453562
agegrp3						
1	.0715136	.0253373	2.82	0.005	.0215432	.121484
2	.130465	.0280521	4.65	0.000	.0751404	.1857895
lnBMI25	.2898622	.0772432	3.75	0.000	.1375229	.4422015
_cons	4.789641	.0224814	213.05	0.000	4.745303	4.833979

Note 0 is missing: it is base/reference group

Morten Frydenberg Linear and Logistic regression - Note 2.2 5

Categorical variable : age group 0 reference

$$\ln(sbp) = \alpha_0 + \alpha_1 \cdot age1 + \alpha_2 \cdot age2 + \alpha_3 \cdot woman + \alpha_4 \cdot \ln\left(\frac{bmi}{25}\right) + E$$

Adjusted difference between for a person in age group 1 compared to age group 0

lnSBP	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
woman	.0035403	.0212026	0.17	0.868	-.0382757	.0453562
agegrp3						
1	.0715136	.0253373	2.82	0.005	.0215432	.121484
2	.130465	.0280521	4.65	0.000	.0751404	.1857895
lnBMI25	.2898622	.0772432	3.75	0.000	.1375229	.4422015
_cons	4.789641	.0224814	213.05	0.000	4.745303	4.833979

Adjusted difference between for a person in age group 2 compared to age group 0

Expected value for a man in age group 0 with bmi=25.

Morten Frydenberg Linear and Logistic regression - Note 2.2 6

The expected values:

$$\text{age} < 40: \ln(\text{sbp}) = \alpha_0 + \alpha_3 \cdot \text{woman} + \alpha_4 \cdot \ln(\text{bmi}/25)$$
$$40 \leq \text{age} < 50: \ln(\text{sbp}) = \alpha_0 + \alpha_1 + \alpha_3 \cdot \text{woman} + \alpha_4 \cdot \ln(\text{bmi}/25)$$
$$50 \leq \text{age}: \ln(\text{sbp}) = \alpha_0 + \alpha_2 + \alpha_3 \cdot \text{woman} + \alpha_4 \cdot \ln(\text{bmi}/25)$$

The estimates

	lnSBP	Coef.	
woman	.003540	3	
1. agegrp3	.071514	1	
2. agegrp3	.130465	2	
lnbmi25	.289862	4	
_cons	4.789641	0	

$$\text{age} < 40: \ln(\text{sbp}) = 4.789 + 0.004 \cdot \text{woman} + 0.290 \cdot \ln(\text{bmi}/25)$$
$$40 \leq \text{age} < 50: \ln(\text{sbp}) = 4.789 + 0.072 + 0.004 \cdot \text{woman} + 0.290 \cdot \ln(\text{bmi}/25)$$
$$50 \leq \text{age}: \ln(\text{sbp}) = 4.789 + 0.130 + 0.004 \cdot \text{woman} + 0.290 \cdot \ln(\text{bmi}/25)$$


Morten FrydenbergLinear and Logistic regression - Note 2.27

Plot02

Agegroup

	Women	Men
0:	$4.793 + 0.290 \cdot \ln(\text{bmi}/25)$	$4.790 + 0.290 \cdot \ln(\text{bmi}/25)$
1:	$4.865 + 0.290 \cdot \ln(\text{bmi}/25)$	$4.861 + 0.290 \cdot \ln(\text{bmi}/25)$
2:	$4.924 + 0.290 \cdot \ln(\text{bmi}/25)$	$4.920 + 0.290 \cdot \ln(\text{bmi}/25)$

Morten FrydenbergLinear and Logistic regression - Note 2.28



Morten FrydenbergLinear and Logistic regression - Note 2.29

Categorical variable : age group 1 reference

$$\ln(\text{sbp}) = \gamma_0 + \gamma_1 \cdot \text{age0} + \gamma_2 \cdot \text{age2} + \gamma_3 \cdot \text{woman} + \gamma_4 \cdot \ln\left(\frac{\text{bmi}}{25}\right) + E$$

age0 is one if a person is in age group 0 and zero otherwise
 age2 is one if a person is in age group 2 and zero otherwise

The expected $\ln(\text{sbp})$ in the three age groups will be:

$$\text{age} < 40: \ln(\text{sbp}) = \gamma_0 + \gamma_1 + \gamma_3 \cdot \text{woman} + \gamma_4 \cdot \ln(\text{bmi}/25)$$
$$40 \leq \text{age} < 50: \ln(\text{sbp}) = \gamma_0 + \gamma_3 \cdot \text{woman} + \gamma_4 \cdot \ln(\text{bmi}/25)$$
$$50 \leq \text{age}: \ln(\text{sbp}) = \gamma_0 + \gamma_2 + \gamma_3 \cdot \text{woman} + \gamma_4 \cdot \ln(\text{bmi}/25)$$

We see that γ_1 is the adjusted difference in $\ln(\text{sbp})$ when comparing a person in the **first group** with one in the **second group**.
And γ_2 is the adjusted difference in $\ln(\text{sbp})$ when comparing a person in the **third group** with one in the **second group**.

Morten FrydenbergLinear and Logistic regression - Note 2.210

Categorical variable : age group 1 reference

$$\ln(\text{sbp}) = \gamma_0 + \gamma_1 \cdot \text{age0} + \gamma_2 \cdot \text{age2} + \gamma_3 \cdot \text{woman} + \gamma_4 \cdot \ln\left(\frac{\text{bmi}}{25}\right) + E$$

Finally we see that γ_0 is the expected $\ln(\text{sbp})$ for a man in the **second** (reference) age group, with $\text{bmi}=25$.

Many programs (but regression in SPSS) let you chose the reference group
In Stata 11 this done like this:

```
regress lnSBP woman b1.agegrp3 lnbmi25
```

This is categorical with base/reference set to 1

Morten FrydenbergLinear and Logistic regression - Note 2.211

Categorical variable : age group 1 reference

$$\ln(\text{sbp}) = \gamma_0 + \gamma_1 \cdot \text{age0} + \gamma_2 \cdot \text{age2} + \gamma_3 \cdot \text{woman} + \gamma_4 \cdot \ln\left(\frac{\text{bmi}}{25}\right) + E$$

Source	SS	df	MS	Number of obs =
Model	.980169926	4	.245042482	200
Residual	4.26524771	195	.021873065	F(4, 195) = 11.20
Total	5.24541764	199	.026358883	Prob > F = 0.0000
				R-squared = 0.1869
				Adj R-squared = 0.1702
				Root MSE = .1479

	lnSBP	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]
woman		.0035403	.0212026	0.17	0.868	-.0382757 .0453562
agegrp3						
0		-.0715136	.0253373	-2.82	0.005	-.121484 -.0215432
2		.0589513	.0263496	2.24	0.026	.0069846 .1109181
lnbmi25		.2898622	.0772432	3.75	0.000	.1375229 .4422015
_cons		4.861154	.0207406	234.38	0.000	4.82025 4.902059

Note 1 is missing: it is base/reference group

Morten FrydenbergLinear and Logistic regression - Note 2.212

Categorical variable : age group 1 reference

$$\ln(sbp) = \gamma_0 + \gamma_1 \cdot age0 + \gamma_2 \cdot age2 + \gamma_3 \cdot woman + \gamma_4 \cdot \ln\left(\frac{bmi}{25}\right) + E$$

Adjusted difference between for a person in age group 0 compared to age group 1

	lnSBP	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]
woman		.0035403	.0212026	0.17	0.868	-.0382757 .0453562
agegrp3						
0		-.0715136	.0253373	-2.82	0.005	-.121484 -.0215432
2		-.0589513	.0263496	-2.24	0.026	-.0069846 .1109181
lnBMI25		.2898622	.0772432	3.75	0.000	.1375229 .4422015
_cons		4.861154	.0207406	234.38	0.000	4.82025 4.902059

Adjusted difference between for a person in age group 2 compared to age group 1

Expected value for a man in age group 1 with bmi=25.

Morten FrydenbergLinear and Logistic regression - Note 2.213

Categorical variable: Comparing to parameterizations

$$\ln(sbp) = \alpha_0 + \alpha_1 \cdot age1 + \alpha_2 \cdot age2 + \alpha_3 \cdot woman + \alpha_4 \cdot \ln\left(\frac{bmi}{25}\right) + E$$

$$\ln(sbp) = \gamma_0 + \gamma_1 \cdot age0 + \gamma_2 \cdot age2 + \gamma_3 \cdot woman + \gamma_4 \cdot \ln\left(\frac{bmi}{25}\right) + E$$

age group 0: $\alpha_0 = \gamma_0 + \gamma_1$

age group 1: $\alpha_0 + \alpha_1 = \gamma_0$

age group 2: $\alpha_0 + \alpha_2 = \gamma_0 + \gamma_2$

$$\alpha_0 = \gamma_0 + \gamma_1$$

$$\alpha_1 = -\gamma_1$$

$$\alpha_2 = \gamma_2 - \gamma_1$$

$$\alpha_3 = \gamma_3$$

$$\alpha_4 = \gamma_4$$

$$\gamma_0 = \alpha_0 + \alpha_1$$

$$\gamma_1 = -\alpha_1$$

$$\gamma_2 = \alpha_2 - \alpha_1$$

$$\gamma_3 = \alpha_3$$

$$\gamma_4 = \alpha_4$$

Morten FrydenbergLinear and Logistic regression - Note 2.214

Categorical variable: Comparing two parameterizations

The estimates:

age group 0 reference

age group 1 reference

Root MSE	=	.1479	
	lnSBP	Coef.	[95% CI]
woman		.0035	-.0382 .0453
agegrp3			
1		.0715	.0215 .1214
2		.1304	.0751 .1857
lnBMI25		.2898	.1375 .4422
_cons		4.7896	4.745 4.833

Root MSE	=	.1479	
	lnSBP	Coef.	[95% CI]
woman		.0035	-.0382 .0453
agegrp3			
0		-.0715	-.1214 -.0215
2		.0589	.0069 .1109
lnBMI25		.2898	.1375 .4422
_cons		4.8611	4.820 4.902

Note, the estimates fulfil the same equations.

The interpretation of the "agegrp3 2 line" and "_cons line" are altered!!!!!!!!

Always remember: what is the reference group!

Morten FrydenbergLinear and Logistic regression - Note 2.215

Categorical variable : age group 1 reference

	lnSBP	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]
woman		.0035403	.0212026	0.17	0.868	-.0382757 .0453562
agegrp3						
0		-.0715136	.0253373	-2.82	0.005	-.121484 -.0215432
2		-.0589513	.0263496	-2.24	0.026	-.0069846 .1109181
lnBMI25		.2898622	.0772432	3.75	0.000	.1375229 .4422015
_cons		4.861154	.0207406	234.38	0.000	4.82025 4.902059

Two test:

One testing no difference between age group 0 and 1.

One testing no difference between age group 2 and 1.

Can we get one test testing no difference between age groups?

A F-test in Stata: `testparm i.agegrp`

(1) 0.agegrp3 = 0

(2) 2.agegrp3 = 0

F(2, 195) = 10.93

Prob > F = 0.0000

Highly significant

Morten FrydenbergLinear and Logistic regression - Note 2.216

Interactions/effectmodification

$$\ln(sbp) = \gamma_0 + \gamma_1 \cdot age0 + \gamma_2 \cdot age2 + \gamma_3 \cdot woman + \gamma_4 \cdot \ln\left(\frac{bmi}{25}\right) + E$$

One of the central assumptions was "no effect modification".

E.g. in model above the "effect" of age, sex and bmi did not depend on the value of each other.

One can introduce effect modification between a categorical variable and another variable.

Here we first will look at agegrp3 and lnBMI25.

The effect modification will be that the coefficient to lnBMI25 depend on age group.

That is, we will allow different effect of bmi in the different age groups.

Morten FrydenbergLinear and Logistic regression - Note 2.217

Interactions/effectmodification

$$\ln(sbp) = \omega_0 + \omega_1 \cdot age0 + \omega_2 \cdot age2 + \omega_3 \cdot woman$$

$$+ \omega_4 \cdot \ln\left(\frac{bmi}{25}\right) + \omega_5 \cdot age0 \cdot \ln\left(\frac{bmi}{25}\right) + \omega_6 \cdot age2 \cdot \ln\left(\frac{bmi}{25}\right) + E$$

$$age \leq 40: \ln(sbp) = (\omega_0 + \omega_1) + (\omega_4 + \omega_5) \cdot \ln\left(\frac{bmi}{25}\right) + \omega_3 \cdot woman$$

$$40 < age \leq 50: \ln(sbp) = \omega_0 + \omega_4 \cdot \ln\left(\frac{bmi}{25}\right) + \omega_3 \cdot woman$$

$$50 < age: \ln(sbp) = (\omega_0 + \omega_2) + (\omega_4 + \omega_6) \cdot \ln\left(\frac{bmi}{25}\right) + \omega_3 \cdot woman$$

$$\omega_1$$
 is the difference between the constant for age group 0 and reference group.

$$\omega_5$$
 is the difference between the coefficient to lnBMI25 for age group 0 and reference group.

Morten FrydenbergLinear and Logistic regression - Note 2.218

Interactions/effectmodification					
regress	lnSBP	woman	b1.agegrp3#c.lnBMI25		
Source	SS	df	MS	Number of obs = 200	
Model	.994860827	6	.165810138	F(6, 193) = 7.53	
Residual	4.25055681	193	.02202361	Prob > F = 0.0000	
Total	5.24541764	199	.026358883	R-squared = 0.1897	
				Adj R-squared = 0.1645	
				Root MSE = .1484	
lnSBP	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]
woman	-.0076438	.0219199	0.35	0.728	-.0355896 .0508772
agegrp3					
0	-.0708045	.0261198	-2.71	0.007	-.1223213 -.0192877
2	.0631082	.0270342	2.33	0.021	.0097877 .1164287
lnBMI25	.3155479	.1222905	2.58	0.011	.0743505 .5567453
agegrp3#c.lnBMI25					
0	-.0429736	.1912373	0.22	0.822	-.3342099 .420157
2	-.1165375	.1855477	-0.63	0.531	-.4824991 .2494242
_cons	4.859743	.0214122	226.96	0.000	4.817511 4.901975


Morten FrydenbergLinear and Logistic regression - Note 2.219

Interactions/effect modification					
Ref: constant and 'slope' in reference group					
0: difference in constant and slope compared to reference					
2: difference in constant and slope compared to reference					
lnSBP	Coef.	Std. E.	t	P> t	[95% Conf. Interval]
woman	-.0076438	.0219199	0.35	0.728	-.0355896 .0508772
agegrp3					
0	-.0708045	.0261198	-2.71	0.007	-.1223213 -.0192877
2	.0631082	.0270342	2.33	0.021	.0097877 .1164287
lnBMI25	.3155479	.1222905	2.58	0.011	.0743505 .5567453
agegrp3#c.lnBMI25					
0	-.0429736	.1912373	0.22	0.822	-.3342099 .420157
2	-.1165375	.1855477	-0.63	0.531	-.4824991 .2494242
_cons	4.859743	.0214122	226.96	0.000	4.817511 4.901975

Note the larger standard errors

Based on the estimates one can find the six "dose-response" curves:

Morten FrydenbergLinear and Logistic regression - Note 2.220



Interactions/effect modification					
lnSBP	Coef.	Std. E.	t	P> t	[95% Conf. Interval]
woman	-.0076438	.0219199	0.35	0.728	-.0355896 .0508772
agegrp3					
0	-.0708045	.0261198	-2.71	0.007	-.1223213 -.0192877
2	.0631082	.0270342	2.33	0.021	.0097877 .1164287
lnBMI25	.3155479	.1222905	2.58	0.011	.0743505 .5567453
agegrp3#c.lnBMI25					
0	-.0429736	.1912373	0.22	0.822	-.3342099 .420157
2	-.1165375	.1855477	-0.63	0.531	-.4824991 .2494242
_cons	4.859743	.0214122	226.96	0.000	4.817511 4.901975

Two tests:

One testing differences between "slope" in age group 0 and 1.

One testing differences between "slope" in age group 2 and 1.

One test testing no difference between age groups!

A F-test in Stata: `testparm i.agegrp3#c.lnBMI25`

(1) 0.agegrp3#c.lnBMI25 = 0

(2) 2.agegrp3#c.lnBMI25 = 0

F(2, 193) = 0.33

Prob > F = 0.7168

Non-significant

Morten FrydenbergLinear and Logistic regression - Note 2.222

Interactions/effect modification					
The test of no interaction was non-significant.					
But look at the confidence interval for the difference in slope for between age group 2 and group 1!					
lnSBP	Coef.	Std. E.	t	P> t	[95% Conf. Interval]
woman	-.0076438	.0219199	0.35	0.728	-.0355896 .0508772
agegrp3					
0	-.0708045	.0261198	-2.71	0.007	-.1223213 -.0192877
2	.0631082	.0270342	2.33	0.021	.0097877 .1164287
lnBMI25	.3155479	.1222905	2.58	0.011	.0743505 .5567453
agegrp3#c.lnBMI25					
0	-.0429736	.1912373	0.22	0.822	-.3342099 .420157
2	-.1165375	.1855477	-0.63	0.531	-.4824991 .2494242
_cons	4.859743	.0214122	226.96	0.000	4.817511 4.901975

It is very wide!!! We know very little about this difference!

The test for no interaction has very low power!!!

The data have very little information on whether there is effect modification.

Morten FrydenbergLinear and Logistic regression - Note 2.223

Interaction between age group and sex					
regress	lnSBP	lnBMI25	b1.agegrp3##b1.sex	Male sex==1	
lnSBP	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]
lnBMI25	.2265018	.0774898	2.92	0.004	.0736662 .3793374
agegrp3					
0	-.0426734	.0377096	-1.13	0.259	-.1170493 .0317025
2	-.0025412	.0365457	-0.07	0.945	-.0746215 .0695391
2.sex	-.0210869	.0322283	-0.65	0.514	-.0846518 .042478
agegrp3#sex					
0 2	-.0548967	.0501668	-1.09	0.275	-.1538422 .0440488
2 2	.133379	.0501308	2.66	0.008	.0345043 .2322536
_cons	4.873442	.0251767	193.57	0.000	4.823786 4.923099

testparm i.agegrp3#i.sex

(1) 0.agegrp3#2.sex = 0

(2) 2.agegrp3#2.sex = 0

F(2, 193) = 6.26

Prob > F = 0.0023

Highly significant

Morten FrydenbergLinear and Logistic regression - Note 2.224

Interaction between age group and sex

Differences between age groups among men are small

	lnSBP	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]
lnBMI25		.2265018	.0774898	2.92	0.004	.0736662 .3793374
agegrp3						
0		-.0426734	.0377096	-1.13	0.259	-.1170493 .0317025
2		-.0025412	.0365457	-0.07	0.945	-.0746215 .0695391
2.sex		-.0210869	.0322283	-0.65	0.514	-.0846518 .042478
agegrp3#sex						
0 2		-.0548967	.0501668	-1.09	0.275	-.1538422 .0440488
2 2		.133379	.0501308	2.66	0.008	.0345043 .2322536
_cons		4.873442	.0251767	193.57	0.000	4.823786 4.923099

Women age group 1: $4.873 - 0.021 = 4.852$
Women age group 0: $4.873 - 0.021 - 0.042 = 4.755$
Women age group 2: $4.873 - 0.021 - 0.003 + 0.133 = 4.982$

Large differences in the age groups among women.

Morten Frydenberg Linear and Logistic regression - Note 2.2 25

Interaction between age group and sex

Using women as reference: b2 .sex

Large differences between age groups among women

	lnSBP	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]
lnBMI25		.2265018	.0774898	2.92	0.004	.0736662 .3793374
agegrp3						
0		-.0975701	.0328469	-2.97	0.003	-.162355 -.0327852
2		.1308378	.0354804	3.69	0.000	.0608587 .2008168
1.sex		.0210869	.0322283	0.65	0.514	-.042478 .0846518
agegrp3#sex						
0 1		.0548967	.0501668	1.09	0.275	-.0440488 .1538422
2 1		-.133379	.0501308	-2.66	0.008	-.2322536 -.0345043
_cons		4.852355	.0205502	236.12	0.000	4.811824 4.892887

Morten Frydenberg Linear and Logistic regression - Note 2.2 26

Plot05

The graph shows six lines representing different combinations of sex and age group. For women, the lines are solid (40), dotted (40-50), and dashed (50+). For men, the lines are dashed (40), dotted (40-50), and solid (50+). All lines show a positive correlation between BMI and expected ln(SBP). The lines for men are generally higher than those for women, and the lines for older age groups are higher than those for younger age groups within each sex.

Agegroup Women Men

0: $4.755 + 0.227 \cdot \ln(bmi/25)$ $4.831 + 0.227 \cdot \ln(bmi/25)$

1: $4.852 + 0.227 \cdot \ln(bmi/25)$ $4.873 + 0.227 \cdot \ln(bmi/25)$

2: $4.982 + 0.227 \cdot \ln(bmi/25)$ $4.871 + 0.227 \cdot \ln(bmi/25)$

Morten Frydenberg Linear and Logistic regression - Note 2.2 27